


Rethinking peatland fire: Fire history of peatlands in the Great Lakes Region



Colleen M. Sutheimer
SILVIS Lab, Forest and Wildlife Ecology
University of Wisconsin-Madison





What do three of the largest wildfires in recent Great Lakes history have in common?


Hiawatha NF/USFS

Where did this fire start?



Mixed messages in mixed landscapes





Objectives

- Determine frequency of low- and moderate- severity fire in Great Lakes' peatlands using tree-rings
- Understand drivers of historical peatland fire regimes
- Compare fire regimes among tree-rings, peat sediments, and lake sediments

Goals


- Characterize historical Great Lakes' peatland fire regimes
- Inform fire management, fire models, and fire risk assessments

Fire scars and tree-rings


- Identify individual fire events including year and season
- Capture widespread low- and moderate-severity fire
- Record 100s to 1000s years of fire history



	No. samples	Area sampled (ha)	Minimum Mean Fire Return Interval (yr)	Maximum Mean Fire Return Interval (yr)	Time Span
Haymeadow Flowage (a)	26	82	7	18	1697–2019
Ramsey Lake (b)	41	278	12	25	1570–2018
Betchler Lake (c)	62	1217	9	34	1520–2018




Haymeadow Flowage(a)




Widespread fire years:
1845, 1847, 1860, 1863,
1866, 1868, 1874, and 1891

Ramsey Lake(b)







Widespread fire years:
1665, 1681, 1702, 1755,
1791, 1792, 1869, 1887,
1900, 1907, 1920, and 1934

Betchler Lake(c)




Widespread fire years:
1718, 1733, 1737, 1751,
1754, 1774, 1847, 1891, and
1920

Widespread peatland fires occurred during dry conditions but not severe drought



Peatland fires occurred in late summer to early spring


	% Dormant	% Earlywood	% Latewood
Haymeadow Flowage	88.5	11.5	0
Ramsey Lake	23.6	8.3	68.1
Betchler Lake	80.8	0	18.2




Charcoal and sediments

- Identify fire periods not fire events
- Date using radiocarbon and age-depth modelling
- Record 100s to 1000s of years of fire history
- Charcoal type linked to fuel type









Fine fuels burned during fire frequent periods at Betchler Lake


Frothy(Forbs, Pine needles and twigs, grass)



Grass (Big blue stem, Pine twigs, Indian grass)



Fibrous (Pine needles, forbs, grass, Pine wood)



Rethinking peatland fire in the Great Lakes

- ❑ **Convention:** Fires only burned every 100 to 1000 years, required severe sustained drought, burn from uplands into peatlands
- ❑ **Our findings:** Fires burned every 10 years, burned under dry conditions but not drought, widespread across uplands and peatlands
- ❑ **Fire management in peatlands now, determines resilience and conservation of peatlands in future**

Acknowledgements

Committee Members: Dr. Volker Radeloff, Dr. Sara Hotchkiss, and Dr. Jed Meunier

Wisconsin DNR Office of Applied Science, Wildlife Management, Forestry and Natural Heritage Conservation Divisions

WDNR: Abe Lenocho, Megan Sebasky, Matt Heritsch, Ivy Widick, and Dr. James Riser

UW-Madison: Cassandra Kaplon

Hiawatha NF: Eric Rebitzke, Brenda Dale, Sam Barnes, Sam Duerksen, and Cory Henry

Ottawa NF: Forest Paukert and Jeremy Erickson

Chequamegon Nicolet NF: John Lampereur, Jay Saunders(Ret.), Scott Linn, and Tym Sauter

Michigan Technological University: Dominic Uhelski, Dr. Evan Kane, and Dr. Rodney Chimner

Michigan DNR Forest Resources Division: Keith Magnusson

Michigan Natural Features Inventory: Josh Cohen

The Nature Conservancy: Randy Swaty

